

Scale ratio ICP for 3D registration of coronary venous anatomy with left ventricular epicardial surface to guide CRT lead placement (Erratum)

Haipeng Tang^a, Robert Bober^b, Chaoyang Zhang^a, Zhuo He^a, Jiangang Zou^c, and Weihua Zhou^a

^aThe Univ. of Southern Mississippi (United States)

^bOchsner Medical Ctr. (United States)

^cFirst Affiliated Hospital of Nanjing Medical Univ. (China)

[Proceedings Volume 10951, Medical Imaging 2019: Image-Guided Procedures, Robotic Interventions, and Modeling; 1095123 \(2019\) <https://doi.org/10.1117/12.2512775>](#)

Event: SPIE Medical Imaging, 2019, San Diego, California, United States

Online Publication Date: 8 March 2019

Erratum Published: 14 June 2019

A revised version of this manuscript was published on 14 June 2019. Details of the revision are provided below. The original paper has been updated.

1, in Eq.2, a transpose symbol is added to the second term of the equation. Before $2\sum_{i=1}^{Np} S \cdot R\tilde{p}_i\tilde{m}_i^k$, after $2\sum_{i=1}^{Np} S \cdot (R\tilde{p}_i)^{(T)}\tilde{m}_i^k$.

2, in Eq.3, a minus symbol is added to the matrix. Before $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$, after $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$.

3, in Eq.4, a transpose symbol is added to the numerator. Before $\sum_{i=1}^{Np} R^{k+1}\tilde{p}_i\tilde{m}_i^k$, after $\sum_{i=1}^{Np} (R^{k+1}\tilde{p}_i)^{(T)}\tilde{m}_i^k$.